

**Queen's University**  
**Faculty of Arts and Science**  
**Department of Mathematics and Statistics**

MTHE 224 – C. Kennedy  
December 2024

<b>HAND IN</b> Answers recorded on exam paper
--

- You have 3 hours for this final exam.
- No aids other than your calculator (Casio 991 series) are allowed.
- There are 7 questions worth a total of 80 points.
- For full marks, you must show all your work and explain how you arrived at your answers unless explicitly told to do otherwise.
- Proctors cannot respond to queries about the interpretation of exam questions. Do your best to answer exam questions as written.

This material is copyrighted solely for students registered in **MTHE 224** and writing this exam. This material shall not be distributed or disseminated. Failure to abide by these conditions is a breach of copyright and may constitute a breach of academic integrity under the University Senate's Academic Integrity Policy Statement.

Do not start the test until your invigilator instructs you to do so.

## Problem 1

(10pts) Consider the ordinary differential equation

$$y'(x) = \frac{x}{y} - xy. \quad (1)$$

(a) Substituting  $z = y^2(x)$ , show that

$$z' + 2xz = 2x. \quad (2)$$

(b) Hence, find the solution to Equation (1) on  $(0, \infty)$  when  $y(0) = 2$ .

*Continue your solution of Problem 1 here.*

## Problem 2

(10pts) Consider the ordinary differential equation

$$y'(x) = f\left(\frac{y}{x}\right), \quad x > 0. \quad (3)$$

(a) Substituting  $y = xv(x)$ , show that

$$\int \frac{dv}{f(v) - v} = \ln(x) + C. \quad (4)$$

(b) Hence, find the solution for  $f(v) := v^2 + 3v + 1$  when  $y(1) = 1$ .

*Continue your solution of Problem 2 here.*

### Problem 3

(10pts) Consider the ordinary differential equation

$$3y^3e^{3xy} - 1 + (2ye^{3xy} + 3xy^2e^{3xy})y'(x) = 0. \quad (5)$$

- (a) Show that Equation (5) is exact.
- (b) Hence, find the solution to Equation (5) when  $y(0) = 1$ .

*Continue your solution of Problem 3 here.*

## Problem 4

(10pts) Consider functions  $\phi_1(t) = t^3$  and  $\phi_2(t) = t^2|t|$ .

- (a) Show that  $\{\phi_1(t), \phi_2(t)\}$  is a linearly dependent set on  $(0, \infty)$ .
- (b) Show that  $\{\phi_1(t), \phi_2(t)\}$  is linearly independent set on  $(-\infty, \infty)$ .



*Continue your solution of Problem 4 here.*

## Problem 5

(10pts) Consider the ordinary differential equation

$$y'' - 4y' - 12y = 36e^{4t}. \quad (6)$$

(a) Find the general solution to

$$y'' - 4y' - 12y = 0. \quad (7)$$

(b) Find the general solution to Equation (6). (*Hint: Use undetermined coefficients.*)

*Continue your solution of Problem 5 here.*

## Problem 6

(15pts) Consider the ordinary differential equation

$$ty'' - (t + 1)y' + y = t^2. \quad (8)$$

(a) Without solving Equation (8), find the Wronskian  $W$ . (*Hint: Use Abel's identity.*)

(b) Show that  $y_1(t) = e^t$  solves the homogeneous equation

$$ty'' - (t + 1)y' + y = 0. \quad (9)$$


(c) Find the general solution to Equation (9). (*Hint: Use the Wronskian.*)

(d) Find the general solution to Equation (8). (*Hint: Use variation of parameters.*)

*Continue your solution of Problem 6 here.*

## Problem 7

(15pts) Suppose a cantilevered beam has length  $L$  [m], a uniform load  $w$  [N]  $\cdot$  [m] $^{-1}$  and constant flexural rigidity  $EI$  [N]  $\cdot$  [m] $^2$ .



The differential equation of the deflection curve is

$$EIy^{(4)}(x) = -w, \quad (10)$$

where  $y(x)$  is the height of the beam relative to the fixed end at  $A$ .

(a) Find the general solution to Equation (10).

For simplicity, assume that  $EI = 1$ ,  $L = 1$  and  $w = 24$ . Since the end at  $A$  is fixed,  $y(0) = y'(0) = 0$ . Also, since the end at  $B$  is free,  $y''(L) = y^{(3)}(L) = 0$ . The boundary value problem of the deflection curve is

$$\begin{cases} y^{(4)}(x) = -24 \\ y(0) = y'(0) = 0 \\ y''(1) = y^{(3)}(1) = 0. \end{cases} \quad (11)$$

(b) Solve the boundary value problem (11) for  $y(x)$ .

(c) Find the reaction moment  $M_A$ .

(d) Find the reaction force  $R_A$ .

*Continue your solution of Problem 7 here.*

*Use this page if additional space is required. Clearly state the question number being answered and refer the marker to this page.*



*Use this page if additional space is required. Clearly state the question number being answered and refer the marker to this page.*

# Useful Formulae

## Calculus

$$\begin{aligned}y'(x) &= \lim_{h \rightarrow 0} \frac{y(x+h) - y(x)}{h} \\y(x) &= (g \circ u)(x) \implies y'(x) = g'(u(x)) \cdot u'(x) \\y(x) &= u(x) \cdot v(x) \implies y'(x) = u(x)v'(x) + u'(x)v(x) \\y(x) &= \frac{u(x)}{v(x)} \implies y'(x) = \frac{u'(x)v(x) - u(x)v'(x)}{v(x)^2} \\&\int^x (f \circ g)(x)g'(x) dx = \int^{g(x)} f(u) du \\&\int u dv = uv - \int v du\end{aligned}$$

## First-Order Differential Equations

$$\begin{aligned}y'(x) &= f(x, y) \implies y_{n+1} = y_n + (x_{n+1} - x_n)f(x_n, y_n) \\y'(x) + p(x)y &= q(x) \implies \mu(x) = e^{\int^x p(x') dx'} \\M(x, y) + N(x, y)y'(x) &= 0 \quad \text{exact} \iff M_y = N_x \\\psi(x, y) &= \int^x M dx + \int^y N - \partial_y \left( \int^x M dx \right) dy \equiv C\end{aligned}$$

## Higher-Order Differential Equations

$$\begin{aligned}ay'' + by' + cy &= 0 \implies ar^2 + br + c = 0 \\r_1, r_2 \in \mathbb{R}, r_1 \neq r_2 &\implies y(t) = c_1 e^{r_1 t} + c_2 e^{r_2 t} \\r_1 = r_2 = r &\implies y(t) = (c_1 + c_2 t)e^{rt} \\r_{\pm} = \alpha \pm \beta i &\implies y(t) = (c_1 \cos(\beta t) + c_2 \sin(\beta t))e^{\alpha t} \\W(y_1, y_2) &= \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = y_1 y_2' - y_1' y_2 \\y'' + p(t)y' + q(t)y &= 0 \implies W(y_1, y_2) = e^{-\int^t p(s) ds} \\y &= y_h + y_p \quad g(t) = e^{\lambda t} \implies y_p(t) = At^n e^{\lambda t} \\y &= y_1 \int \frac{-g \cdot y_2}{W} + y_2 \int \frac{g \cdot y_1}{W} \\A &= PDP^{-1} \quad y = \sum c_j e^{\lambda_j t} \vec{v}_j \\y'' &= \frac{M}{EI} \quad V(x) = \frac{dM}{dx}\end{aligned}$$